Issue XXXV: Cohomology and Spectra

Maksym Sokhatskyi¹

¹ National Technical University of Ukraine Igor Sikorsky Kyiv Polytechnical Institute 23 травня 2025 р.

Анотація

This article presents formal definitions and theorems for ordinary and generalized cohomology theories, unstable and stable spectra, and spectral sequences in Abelian categories, including the Serre, Atiyah-Hirzebruch, Leray, Eilenberg-Moore, Hochschild-Serre, Filtered Complex, Chromatic, Adams, and Bockstein spectral sequences. We define slopes, sheets, coordinates, quadrants, complex filtrations, and double complexes. Additionally, we explore the categorical foundations of cohomology theories and spectra, including their relationships to algebra, homological algebra, and stable homotopy theory, through isomorphisms, analogies, and instances.

Зміст

1	Stat	ole Ho	motopy Type Theory 2		
	1.1	Ordina	ary Cohomology Theories		
	1.2	2 Generalized Cohomology Theories			
	1.3				
	1.4				
		1.4.1	Algebraic and Spectral Correspondences		
	1.5	Spectr	al Sequences		
		1.5.1	Serre Spectral Sequence		
		1.5.2	Atiyah-Hirzebruch Spectral Sequence		
		1.5.3	Leray Spectral Sequence		
		1.5.4	Eilenberg-Moore Spectral Sequence		
		1.5.5	Hochschild-Serre Spectral Sequence		
		1.5.6	Spectral Sequence of a Filtered Complex		
		1.5.7	Chromatic Spectral Sequence		
		1.5.8	Adams Spectral Sequence		
		1.5.9	Bockstein Spectral Sequence		

1 Stable Homotopy Type Theory

1.1 Ordinary Cohomology Theories

Definition 1. An ordinary cohomology theory on the category of topological spaces and pairs is a contravariant functor $H^*(-; G)$: Top^{op} \rightarrow GrAb, assigning to each pair (X, A) a sequence of abelian groups $\{H^n(X, A; G)\}_{n \in \mathbb{Z}}$, with coefficient group G, satisfying:

- 1. Homotopy: If $f\simeq g:(X,A)\to (Y,B),$ then $f^*=g^*:H^n(Y,B;G)\to H^n(X,A;G).$
- 2. *Exactness*: For (X, A), there is a long exact sequence: $\dots \to H^n(X, A; G) \to H^n(X; G) \to H^n(A; G) \xrightarrow{\delta} H^{n+1}(X, A; G) \to \dots$
- 3. *Excision*: For $U \subset A$ with $\overline{U} \subset int(A)$, the inclusion $(X \setminus U, A \setminus U) \hookrightarrow (X, A)$ induces isomorphisms $H^n(X, A; G) \cong H^n(X \setminus U, A \setminus U; G)$.
- 4. Additivity: For $X = \bigsqcup X_i$, $H^n(X;G) \cong \bigoplus H^n(X_i;G)$.
- 5. Dimension: For a point pt, $H^{n}(pt;G) = \begin{cases} G & n = 0 \\ 0 & n \neq 0 \end{cases}$.

1.2 Generalized Cohomology Theories

Definition 2. A generalized cohomology theory is a contravariant functor h^* : Top^{op} \rightarrow GrAb, assigning to each pair (X, A) a sequence $\{h^n(X, A)\}_{n \in \mathbb{Z}}$, satisfying:

- 1. Homotopy, Exactness, Excision, and Additivity as in Definition 1.
- 2. Suspension: There is a natural isomorphism $h^n(X, A) \cong h^{n+1}(\Sigma X, \Sigma A)$, where Σ is the reduced suspension.

The groups $h^{n}(pt)$ form a graded ring, the coefficients of h^{*} .

Theorem 1. Every generalized cohomology theory h^* is representable by a spectrum $E = \{E_n, \sigma_n : \Sigma E_n \to E_{n+1}\}$, with $h^n(X) \cong [X, E_n]_*$, where $[-, -]_*$ denotes pointed homotopy classes.

1.3 Unstable and Stable Spectra

Definition 3. A spectrum is a sequence of pointed spaces $\{E_n\}_{n \in I}$, where $I \subseteq \mathbb{Z}$, with structure maps $\sigma_n : \Sigma E_n \to E_{n+1}$. It is:

- Unstable if $I \subseteq \mathbb{Z}_{>0}$.
- *Stable* if $I = \mathbb{Z}$ and each σ_n is a homotopy equivalence.

Theorem 2. For an unstable spectrum E, the functor $X \mapsto [X, E_n]_*$ defines a cohomology theory on spaces of dimension $\leq n$. For a stable spectrum E, the functor $h^n(X) = [X, E_n]_*$ defines a generalized cohomology theory.

1.4 Categorical Interpretation

This section explores the categorical foundations of ordinary and generalized cohomology theories, their associated spectra, and the relationships between algebraic and topological categories through isomorphisms, analogies, and instances. We formalize these structures and highlight their categorical nuances isomorphisms and non-isomorphic relationships, drawing on frameworks like algebra, homological algebra, and stable homotopy theory.

Definition 4. The category of spectra, denoted Spectra, is the category whose objects are stable spectra $\mathsf{E} = \{\mathsf{E}_n, \sigma_n : \Sigma \mathsf{E}_n \to \mathsf{E}_{n+1}\}$, where E_n are pointed spaces and σ_n are homotopy equivalences. Morphisms are collections of maps $\mathsf{f}_n : \mathsf{E}_n \to \mathsf{F}_n$ compatible with structure maps. The stable homotopy category is the localization of Spectra at weak equivalences (maps inducing isomorphisms on homotopy groups).

Definition 5. An ordinary cohomology theory is a functor $H^*(-; G) : \operatorname{Top}^{\operatorname{op}} \to \operatorname{GrAb}$ satisfying the Eilenberg-Steenrod axioms (Definition 1). Categorically, it is represented by the Eilenberg-MacLane spectrum H, where $A \in \operatorname{Ab}$, with $H^n(X; A) \cong [X, H_n]_*$.

Definition 6. A generalized cohomology theory is a functor h^* : Top^{op} \rightarrow GrAb satisfying the axioms of Definition 2. It is representable in Spectra, with $h^n(X) \cong [X, E_n]_*$ for a spectrum E.

Theorem 3 (Brown Representability). Every generalized cohomology theory h^* on Top is representable by a spectrum $E \in$ Spectra, i.e., there exists E such that $h^n(X) \cong [X, E_n]_*$ for all $X \in$ Top.

Theorem 4. The stable homotopy category Spectra is a triangulated category, with distinguished triangles corresponding to cofiber sequences. It is equivalent to the category of spectra localized at weak equivalences.

Theorem 5. The functor $A \mapsto H$ from Ab to Spectra, mapping an abelian group to its Eilenberg-MacLane spectrum, is faithful but not full. The induced functor on ordinary cohomology theories to generalized cohomology theories is an embedding of categories.

1.4.1 Algebraic and Spectral Correspondences

Mathematics is unified through *isomorphisms* (categorical equivalences), *analogies* (functorial similarities), and *instances* (specific subcategories or objects). We present a correspondence table linking Algebra (Ab), Homological Algebra (Ch(\mathbb{Z})), Ordinary Cohomology, K-Theory, Superalgebra, and Stable Spectra (Spectra).

Definition 7 (Isomorphism). An **isomorphism** in a category \mathcal{C} is a morphism $f: A \to B$ with an inverse $g: B \to A$ such that $g \circ f = id_A$ and $f \circ g = id_B$. For categories, an isomorphism is an equivalence, i.e., a functor $F: \mathcal{C} \to \mathcal{D}$ with a quasi-inverse $G: \mathcal{D} \to \mathcal{C}$.

Definition 8 (Analogy). A non-isomorphic **analogy** is a structural similarity between objects or categories, captured by functors that preserve some properties but not all, ensuring no categorical equivalence.

Definition 9 (Instance). An **instance** is a specific object or subcategory within a broader category, embedded via a faithful functor. A column in the table is an instance of another if its structures are special cases of the latter's, maintaining non-isomorphic distinctions from other categories.

Category	Object	Ring	Initial Unit	Operations
Algebra	Abelian group	Ring	Z	\oplus, \otimes
Homological Algebra	Chain complex	dg-ring	$\mathbb{Z}[0]$	\oplus, \otimes
Superalgebra	$\mathbb{Z}/2\mathbb{Z}$ -graded Ab	$\mathbb{Z}/2\mathbb{Z}$ -graded Ring	\mathbb{Z}	\oplus, \otimes
Ordinary Cohomology	Cohomology $H^*(-;A)$	Graded ring	$H^*(-;\mathbb{Z})$	\oplus, \otimes
Complex K-Theory	Graded abelian group	Graded ring	KU	\lor,\land
Real K-Theory	Graded abelian group	Graded ring	KO	\lor,\land
Stable Spectra	Stable spectrum	Ring spectrum	S	\lor,\land

Табл. 1: Algebraic and Spectral Correspondences

- Isomorphisms: Rare, e.g., Ab \cong Mod_Z. Most relationships are non-isomorphic.
- Analogies: The tensor product \otimes in Ab and smash product \wedge in Spectra are analogous, but Ab \neq Spectra due to Spectra's triangulated structure.
- *Instances*: KU, KO, and H are instances of Spectra. Superalgebra is an instance of Ab via the forgetful functor.

Example 1. The functor $A \mapsto A[0]$ embeds Ab into $Ch(\mathbb{Z})$, but $Ch(\mathbb{Z}) \neq Ab$ due to differentials. Similarly, $H : Ab \rightarrow$ Spectra embeds abelian groups as Eilenberg-MacLane spectra, but Spectra's stable phenomena (e.g., suspension equivalences) distinguish it.

Remark 1. Non-isomorphic analogies require careful handling. Conflating \land in Spectra with \otimes in Ab can lead to errors in spectral sequence computations, as \land introduces higher Tor terms.

1.5 Spectral Sequences

Definition 10. A spectral sequence in an Abelian category \mathcal{A} is a collection of objects $\{E_r^{p,q}\}_{r>1,p,q\in\mathbb{Z}}, E_r^{p,q} \in \mathcal{A}$, with differentials:

$$\mathbf{d}_{r}^{p,q}: \mathbf{E}_{r}^{p,q} \to \mathbf{E}_{r}^{p+a_{r},q+b_{r}},$$

such that:

- 1. $\mathbf{d}_{\mathbf{r}} \circ \mathbf{d}_{\mathbf{r}} = \mathbf{0}$.
- 2. $E_{r+1}^{p,q} = H^{p,q}(E_r, d_r) = \ker(d_r^{p,q}) / \operatorname{im}(d_r^{p-a_r,q-b_r}).$
- 3. There exists a graded object $H^n\in \mathcal{A}$ with filtration $F_pH^{p+q}\subseteq H^{p+q},$ such that:

$$E_{\infty}^{p,q} \cong F_{p}H^{p+q}/F_{p-1}H^{p+q}$$

The sequence is *first-quadrant* if $E_r^{p,q} = 0$ for p < 0 or q < 0.

Definition 11. The r-th sheet of a spectral sequence is the collection $\{E_r^{p,q}\}_{p,q}$. The indices (p,q) are coordinates, with p the filtration degree and q the complementary degree, satisfying total degree n = p + q. The slope of $d_r : E_r^{p,q} \to E_r^{p+r,q-r+1}$ is $\frac{-r+1}{r}$.

Definition 12. A filtered complex in $\mathcal{A} = Ab$ is a chain complex (C_*, ∂) with a filtration $\cdots \subseteq F_{p-1}C_n \subseteq F_pC_n \subseteq F_{p+1}C_n \subseteq \cdots$, compatible with ∂ . A double complex is a bigraded object $C_{p,q}$ with differentials $d^h : C_{p,q} \to C_{p-1,q}$, $d^{\nu} : C_{p,q} \to C_{p,q-1}$, satisfying $d^h d^h = d^{\nu} d^{\nu} = d^h d^{\nu} + d^{\nu} d^h = 0$. The total complex is $Tot(C)_n = \bigoplus_{p+q=n} C_{p,q}$.

Theorem 6. A filtered complex (C_*, F_p) induces a spectral sequence with:

$$\mathsf{E}_0^{\mathfrak{p},\mathfrak{q}} = \mathsf{F}_{\mathfrak{p}}C_{\mathfrak{p}+\mathfrak{q}}/\mathsf{F}_{\mathfrak{p}-1}C_{\mathfrak{p}+\mathfrak{q}}, \quad \mathsf{E}_1^{\mathfrak{p},\mathfrak{q}} = \mathsf{H}_{\mathfrak{p}+\mathfrak{q}}(\mathsf{F}_{\mathfrak{p}}C/\mathsf{F}_{\mathfrak{p}-1}C) \implies \mathsf{H}_{\mathfrak{p}+\mathfrak{q}}(C).$$

A double complex $C_{p,q}$ with filtration by p-index induces:

$$\mathsf{E}_1^{p,q} = \mathsf{H}_q^{\nu}(\mathsf{C}_{p,*}), \quad \mathsf{d}_1 = \mathsf{H}(\mathsf{d}^{\mathsf{h}}) \implies \mathsf{H}_{p+q}(\mathrm{Tot}(\mathsf{C})).$$

1.5.1 Serre Spectral Sequence

Theorem 7. For a fibration $F \rightarrow E \rightarrow B$ with B path-connected, there exists a first-quadrant spectral sequence:

$$\mathsf{E}_{2}^{p,q} = \mathsf{H}^{p}(\mathsf{B};\mathsf{H}^{q}(\mathsf{F};\mathbb{Z})) \implies \mathsf{H}^{p+q}(\mathsf{E};\mathbb{Z}),$$

with $d_r: E_r^{p,q} \to E_r^{p+r,q-r+1}$.

1.5.2 Atiyah-Hirzebruch Spectral Sequence

Theorem 8. For a generalized cohomology theory h^* and a CW-complex X, there exists a spectral sequence:

$$E_2^{p,q} = H^p(X; h^q(pt)) \implies h^{p+q}(X),$$

with $d_r: E_r^{p,q} \to E_r^{p+r,q-r+1}$.

1.5.3 Leray Spectral Sequence

Theorem 9. For a continuous map $f: X \to Y$ and a sheaf \mathcal{F} on X, there exists a spectral sequence:

$$\mathsf{E}_{2}^{p,q} = \mathsf{H}^{p}(\mathsf{Y};\mathsf{R}^{q}\mathsf{f}_{*}\mathcal{F}) \implies \mathsf{H}^{p+q}(\mathsf{X};\mathcal{F}),$$

with $d_r: E_r^{p,q} \to E_r^{p+r,q-r+1}$.

1.5.4 Eilenberg-Moore Spectral Sequence

Theorem 10. For a pullback diagram with fibration $F \to E \to B$, there exists a spectral sequence:

$$\mathsf{E}_{2}^{p,q} = \mathrm{Tor}_{\mathsf{H}_{*}(\mathsf{B})}^{p,q}(\mathsf{H}_{*}(\mathsf{F}),\mathsf{R}) \implies \mathsf{H}_{p+q}(\mathsf{F};\mathsf{R}),$$

with $d_r: E_r^{p,q} \to E_r^{p-r,q+r-1}$.

1.5.5 Hochschild-Serre Spectral Sequence

Theorem 11. For a group extension $1 \rightarrow N \rightarrow G \rightarrow Q \rightarrow 1$, there exists a spectral sequence:

$$\mathsf{E}_{2}^{\mathsf{p},\mathsf{q}} = \mathsf{H}^{\mathsf{p}}(\mathsf{Q};\mathsf{H}^{\mathsf{q}}(\mathsf{N};\mathsf{R})) \implies \mathsf{H}^{\mathsf{p}+\mathsf{q}}(\mathsf{G};\mathsf{R}),$$

with $d_r: E_r^{p,q} \to E_r^{p+r,q-r+1}$.

1.5.6 Spectral Sequence of a Filtered Complex

Theorem 12. For a filtered complex (C_*, F_p) , there exists a spectral sequence:

$$\mathsf{E}_1^{p,q} = \mathsf{H}_{p+q}(\mathsf{F}_pC/\mathsf{F}_{p-1}C) \implies \mathsf{H}_{p+q}(C),$$

with $d_r: E_r^{p,q} \to E_r^{p-r,q+r-1}$.

1.5.7 Chromatic Spectral Sequence

Theorem 13. For a spectrum X, there exists a spectral sequence:

$$\mathsf{E}_1^{n,k} = \pi_{n-k}(\mathsf{L}_{\mathsf{K}(k)}X) \implies \pi_{n-k}(X),$$

where $L_{K(k)}X$ is the localization at the k-th Morava K-theory, with $d_r: E_r^{n,k} \to E_r^{n+1,k-r}$.

1.5.8 Adams Spectral Sequence

Theorem 14. For a spectrum X and prime p, there exists a spectral sequence:

$$\mathsf{E}_2^{s,t} = \mathrm{Ext}_A^{s,t}(\mathrm{Hom}_*(X,\mathbb{Z}/p),\mathbb{Z}/p) \implies \pi_{t-s}(X_{(p)}),$$

where A is the Steenrod algebra, with $d_r: E_r^{s,t} \to E_r^{s+r,t+r-1}.$

1.5.9 Bockstein Spectral Sequence

Theorem 15. For a short exact sequence $0 \to R \to R' \to R'' \to 0$ of coefficient rings, there exists a spectral sequence:

$$\mathsf{E}^{\mathsf{p},\mathsf{q}}_1 = \mathsf{H}^{\mathsf{p}+\mathsf{q}}(\mathsf{X};\mathsf{R}'') \implies \mathsf{H}^{\mathsf{p}+\mathsf{q}}(\mathsf{X};\mathsf{R}),$$

with $d_r: E_r^{p,q} \to E_r^{p+1,q-r}$.

Література

- [1] Hatcher, A., Algebraic Topology, Cambridge University Press, 2002.
- [2] Switzer, R., Algebraic Topology Homology and Homotopy, Springer, 1975.
- [3] Kato, G., The Heart of Cohomology, Springer, 2006.
- [4] Adams, J.F., Stable Homotopy and Generalized Homology, University of Chicago Press, 1974.